



Early Journal Content on JSTOR, Free to Anyone in the World


This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.



Rhacomitrium FLETTII.

PLATE V. Fig. 1. Plants showing mode of branching and disposition of leaves when moist. Natural size about 1.5 cm. \times 3. Fig. 2. Leaves \times 10. Figs. 3, 4, 5. Cross sections of a leaf from near base, middle and apex, respectively. Figs. 6, 7, 8. Leaf areolation from near base, middle and apex, respectively.

RHACOMITRIUM FLETTII n. sp.

JOHN M. HOLZINGER.

Stem densely caespitose, radiculose at base, simple or fasciculately branched; color of plants yellowish brown. Leaves divergent, then ascending when moist, lanceolate, margin entire; costa reaching apex; cells at base pellucid, thin-walled, approximately rectangular, about 1×2 , above more isodiametric, thick-walled, with a row of more pellucid, smaller, roundish cells along the margin and this bistratose toward the apex. Entirely sterile.

This plant was collected by Prof. J. B. Flett on Mt. Tacoma (Mt. Rainier), at an altitude of 14,519 feet, near one of the steam jets issuing from a fissure in the side of the rim of the crater, Aug. 10, 1897.

In appearance, color and branching it looks like a diminutive form of *Rhacomitrium ellipticum*, section *Dryptodon*. But it has almost completely lost the very unequal thickenings of the cell walls above the base, has narrower leaves lacking the foldings in that species, has a heavier costa, and has a tendency to a doubling of the small marginal cells, as is shown in Fig. 5 of the accompanying plate.

Winona, Minn.